

AD-A163 705

CONVERGENCE OF CUBIC SPLINE INTERPOLATION WITH THE
NOT-A-KNOT CONDITION(U) WISCONSIN UNIV-MADISON
MATHEMATICS RESEARCH CENTER C DE BOOR OCT 85

1/1


UNCLASSIFIED

MRC-TSR-2876 DRAG29-80-C-0041

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

2

AD-A163 705

MRC Technical Summary Report #2876

CONVERGENCE OF CUBIC SPLINE INTERPOLATION
WITH THE NOT-A-KNOT CONDITION

Carl de Boor

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

October 1985

(Received October 3, 1985)

DTIC
ELECTE
FEB 5 1986
S B D

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

86 2 5

039

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

CONVERGENCE OF CUBIC SPLINE INTERPOLATION
WITH THE NOT-A-KNOT CONDITION

Carl de Boor

Technical Summary Report #2876
October 1985

ABSTRACT

C² squares
It is shown that cubic spline interpolation with the not-a-knot side condition converges to any C^2 -interpoland without any mesh-ratio restriction as the mesh size goes to zero.

AMS (MOS) Subject Classifications: 41A15, 41A05

Key Words: cubic spline, interpolation, not-a-knot, convergence, total positivity

Work Unit Number 3 (Numerical Analysis and Scientific Computing)

SIGNIFICANCE AND EXPLANATION

The cubic spline interpolant to a given function agrees with that function at the knots $t_1 < \dots < t_n$ of the spline; its complete specification requires two additional conditions. Originally, the slope at the end points t_1 and t_n was used, giving the **complete** spline interpolant. But, in the absence of such information, one has to try something else. One proposal is the **not-a-knot** condition [B₃;Ch. IV] in which the third derivative of P_g is made continuous at t_2 and at t_{n-1} . In effect, these next-to-boundary knots are not knots. But, whereas the convergence of complete spline interpolation has been established in various ways, there doesn't seem to be a proof of the convergence of not-a-knot cubic spline interpolation in the literature.

The present report remedies this situation.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

CONVERGENCE OF CUBIC SPLINE INTERPOLATION

WITH THE NOT-A-KNOT CONDITION

Carl de Boor

The cubic spline interpolant P_g to a given function g agrees with g at the knots $t_1 < \dots < t_n$ of the spline; its complete specification requires two additional conditions. Originally, the slope at the end points t_1 and t_n was used, giving the complete spline interpolant. But, in the absence of such information, one has to try something else. One proposal is the not-a-knot condition [B₃; Ch. IV] in which the third derivative of P_g is made continuous at t_2 and at t_{n-1} . In effect, these next-to-boundary knots are not knots. But, whereas the convergence of complete spline interpolation has been established in various ways, there doesn't seem to be a proof of the convergence of not-a-knot cubic spline interpolation in the literature (except for [SV] in which it appears as a special case but is only treated for a uniform mesh).

The argument here follows the standard path (see [B₁], [B₂]): One considers the derived projector P'' obtained from P by the rule

$$(1) \quad P''D^2g := D^2(Pg)$$

and shows it to be bounded in the uniform norm independently of the mesh, i.e., one shows that

$$(2) \quad M := \sup_t \|P''\| < \infty$$

with

$$\|P''\| := \sup_f \|P''f\|/\|f\|$$

and

$$\|f\| := \|f\|_\infty := \sup_{t_1 \leq x \leq t_n} |f(x)|.$$

Since P'' is a linear projector with range

$$S := \mathcal{S}_{2,s},$$

i.e., with range the span of the linear B-splines on the knot sequence

$$s := (s_i)_{i=1}^{m+2} := (t_1, t_1, t_3, t_4, \dots, t_{n-3}, t_{n-2}, t_n, t_n), \text{ hence } m := n-2,$$

this implies that

$$\|D^2(g - Pg)\| = \|D^2g - P^*(D^2g)\| \leq \|1 - P^*\| \operatorname{dist}(D^2g, S) \leq (1+M) \omega_{D^2g}(|s|)$$

with ω_f the modulus of continuity of f (on $[t_1, t_n]$) and

$$|s| := \max_i \Delta s_i.$$

Consequently,

$$\|g - Pg\| \leq ((1+M)|t|^2/8)(|s|^2/8)\|D^4g\|,$$

using the fact that Pg agrees with g at t and using the standard estimate for $\operatorname{dist}(D^2g, S)$ in case $D^2(D^2g)$ is bounded.

This leaves the hard part of the argument, namely a proof of (2), but here, too, the approach is standard, as follows.

Since Pg agrees with g at t , P^*D^2g agrees with D^2g at

$$\lambda_i := \int M_i, \quad i=1, \dots, m,$$

with $M_i := M_{i,2,t}$ the linear B-splines for the knot sequence t (and normalized to have $\int M_i = 1$). This follows from the fact

$$2! [t_i, t_{i+1}, t_{i+2}]g = \int M_i(x) D^2g(x) dx.$$

Hence, with $N_i := N_{i,2,s}$ the linear B-splines for the knot sequence s (and normalized to sum to 1), and $f := D^2g$, we obtain P^*f in the form

$$P^*f =: \sum a_i N_i,$$

where a is the unique solution of the linear system

$$(3) \quad Aa = (\lambda_i f)$$

with

$$(4) \quad A := \left(\int M_i N_j \right).$$

Consequently,

$$\|P^*f\| = \|a\|_\infty \leq \|A^{-1}\|_\infty \max_i |\int M_i f| \leq \|A^{-1}\|_\infty \|f\|_\infty.$$

This shows that

$$(5) \quad \|P^*\| = \|A^{-1}\|_\infty$$

and so reduces the proof of (2) to finding a uniform lower bound for A .

Since A is totally positive, a lower bound for A is obtained from any vector x for which

$$d := \min_j (-)^j (Ax)_j > 0.$$

For, the total positivity of A implies (by Cramer's rule) that its inverse is checkerboard, i.e.,

$$(-)^{i+j} A^{-1}(i,j) > 0.$$

Therefore,

$$|x(i)| = |\sum A^{-1}(i,j) (Ax)_j| = \sum |A^{-1}(i,j)| (-)^j (Ax)_j > \sum |A^{-1}(i,j)| d$$

hence

$$\|x\|_\infty = \max_i |x_i| > \|A^{-1}\|_\infty d$$

or

$$(6) \quad \|A^{-1}\|_\infty < \|x\|_\infty / d.$$

The matrix $3A$ is tridiagonal, with typical row

$$\Delta s_1 / (s_{i+2} - s_i), \quad 2, \quad \Delta s_{i+1} / (s_{i+2} - s_i), \quad i=3, \dots, m-2,$$

and only the first two and the last two rows deviate from this. Here are the first two rows:

$$\begin{aligned} & 1 + (\Delta t_2) / (s_3 - s_1), \quad 2 - (\Delta t_2) / (s_3 - s_1) \\ & \frac{(\Delta t_2)^2}{(s_3 - s_1)(t_4 - t_2)}, \quad \frac{1}{(t_4 - t_2)} \left[\frac{2(\Delta t_2)^2 + 3\Delta t_2 \Delta t_1}{(s_3 - s_1)} + 2\Delta t_3 \right], \quad \Delta s_3 / (t_4 - t_2) \end{aligned}$$

The last two rows can be obtained from this by the appropriate change in variables.

The typical row of $3A$ suggests the vector $x := (-1, 1, -1, 1, \dots)$ as suitable since this gives

$$b := 3Ax = (b_1, b_2, -1, 1, -1, 1, \dots, b_{m-1}, b_m),$$

i.e., an appropriate result except, perhaps, for the first two and last two components.

One can check that, as t_2 varies in the interval $[s_1, s_3]$, the first component of b becomes positive and this destroys the desired alternation. This can be helped, though, by choosing x_1 and x_2 in dependence on t_2 . Specifically, I choose x in the form

$$x = (x_1, x_2, -1, 1, -1, 1, \dots)$$

so that

$$b = (-1, 1, b_3, 1, -1, 1, \dots) ,$$

with the analogous happening at the other end. This requires that $m > 5$, to avoid destructive interference of the machinations on one end with those at the other. With this assumption, though, it is sufficient to run through the details for the left end only. This means that I solve the linear system

$$\begin{aligned} A(1,1)x_1 + A(1,2)x_2 &= -1 \\ A(2,1)x_1 + A(2,2)x_2 &= 1 - A(2,3)^*(-1) \end{aligned}$$

Since the system is invariant under a linear change in the independent variable, I normalize the situation to one where

$$s_1 = 0, s_3 = 1$$

and define

$$k := \Delta t_2, \quad h := \Delta s_3 .$$

This means that I am interested in the range $k \in [0,1], h > 0$. In these terms, the system reads

$$\begin{vmatrix} 1+k & 2-k \\ k^2 & 3k-k^2+2h \end{vmatrix} x = (-1, k + 2h) .$$

Its solution is

$$x = (-5k + 2k^2 - 6h + 2hk, k + 2k^2 + 2h + 2hk)/D ,$$

with

$$D := (3k + 2h + 2hk) > 0 .$$

Now observe that, for $(k,h) \in [0,1] \times \mathbb{R}_+$,

$$x_1 - (-3) = (4k + 2k^2 + 8hk)/D > 0 \text{ with equality only if } k = 0 ,$$

$$-1 - x_1 = 2(1-k)(k + 2h)/D > 0 \text{ with equality only if } k = 1 .$$

Finally,

$$x_2 = 1 - 2k(1-k)/D < 1 , \text{ with equality only if } k = 0 \text{ or } 1 .$$

I conclude that $x_1 \in [-3, -1]$ and $x_2 \in [2/3, 1]$, hence

$$\|x\|_\infty < 3.$$

Also, $b_3 < -1$, since $x_2 < 1$, while $b_i = (-)^i$ otherwise, hence

$$3d := \min_i (-)^i 3b_i > 1.$$

This shows that

$$(7) \quad \|A^{-1}\|_\infty < 9$$

independently of t (as long as $n > 6$).

Remarks. (i) As $t_2 \rightarrow t_1$ and $t_{n-1} \rightarrow t_n$, not-a-knot cubic spline interpolation reduces to complete spline interpolation, hence P^n becomes least-squares approximation from $\mathcal{S}_{2,t}$. The above argument then reduces to the standard one. In particular, the computed x is just the alternating vector $(-1, 1, -1, 1, \dots)$ and the resulting bound is the customary one: $\|A^{-1}\|_\infty = 3$.

(ii) It may be possible to carry out the argument by perturbation, starting off with the known stability of P^n for complete cubic spline interpolation and showing that the change in the side conditions to the not-a-knot conditions is gentle enough (at least for large n) to change $\|P^n\|$ by a bounded amount.

(iii) When I wrote this note in August of 1984, I wrote that "It should be possible to show the boundedness, independently of t , of the derived linear projector P' , given by the rule

$$P'(Dg) := D(Pg).$$

But the argument via total positivity would be a bit messier." In the meantime, R. K. Beatson, who had earlier brought my attention to the fact that there did not seem to be a convergence proof for spline interpolation with the not-a-knot condition, has independently studied this question and shown that, strictly speaking, P' is not boundable independent of the mesh. The difficulty lies in the first and last data interval. If one measures the size of $P'f$ by the number

$$\|P'f\| := \sup_{t_2 < t < t_{n-1}} |P'f(t)|,$$

then he obtains a mesh-independent bound for $\|P'\| := \sup_f \|P'f'\|/\|f\|$. See Beatson's forthcoming paper "On the convergence of some cubic spline interpolation schemes", ms., Feb. 85, Mathematics, University of Connecticut, Storrs, CN 06268.

REFERENCES

- [B₁] C. de Boor, Bounding the error in spline interpolation, SIAM Review 16 (1974) 531-544.
- [B₂] C. de Boor, On bounding spline interpolation, J. Approximation Theory 14 (1975) 191-203.
- [B₃] C. de Boor, A practical guide to splines, Springer, 1978.
- [SV] B. Swartz & R. S. Varga, Error bounds ofr spline and L-spline interpolation, J. Approximation Theory 6 (1972) 6-49.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2876	2. GOVT ACCESSION NO. ADA 163 705	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CONVERGENCE OF CUBIC SPLINE INTERPOLATION WITH THE NOT-A-KNOT CONDITION		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Carl de Boor		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53705		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis and Scientific Computing
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE October 1985
		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) cubic spline interpolation not-a-knot convergence total positivity		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown that cubic spline interpolation with the not-a-knot side condition converges to any C^2 -interpoland without any mesh-ratio restriction as the size goes to zero.		

END

FILMED

3-86

DTIC